



NOT PREGNANT



PREGNANT



Make Sure You Are Pregnant

Take a urine-based pregnancy test to check if you are pregnant

Check Your Gestation Date

Track the time elapsed since the first day of your last period or use a gestational age calculator.



Check The Abortion Pills

The first pill is MIFEPRISTONE. The second pill to be taken is MISOPROSTOL.



TIMELINE OF PILL USAGE

TIME SINCE LAST PERIOD →	LESS THAN 9 WEEKS	9 TO 12 WEEKS	MORE THAN 12 WEEKS
DAY 1	TAKE MIFEPRISTONE	TAKE MIFEPRISTONE	TAKE MIFEPRISTONE
DAY 2	TAKE 4 MISOPROSTOLS	TAKE 4 MISOPROSTOLS ↓ (4 hrs later) TAKE 4 MORE MISOPROSTOLS ↓ TAKE PAINKILLERS IF REQUIRED	TAKE 2 MISOPROSTOLS EVERY 03 hrs UNTIL PREGNANCY PASSES ↓ PAINKILLERS IF REQUIRED

DAY 1: Take Mifepristone

Take one 200 mg pill of Mifepristone with water



DAY 2: Take Misoprostol

Put the misoprostol pills, as prescribed in the table above under the tongue or inside the vagina.



What to Expect

- Bleeding and cramping start 4 hours or later after taking misoprostol, alongside nausea, vomiting, diarrhea, fever, and chills. These symptoms resolve on their own in a few hours.
- Eating a small amount of food can alleviate nausea.
- Taking misoprostol vaginally may also provide relief



What to Watch Out For

Fever above 102°F, or over 100.4°F for more than 24 hours. Bleeding that fills more than 2 maxi pads. Persistent pain that won't subside despite taking painkillers



Note

- **Less than 9 weeks: Take 04 misoprostol pills.**
- **Between 9 to 12 weeks: Take 04 misoprostol pills. After 4 hours, take 4 more misoprostol pills.**
- **Above 12 weeks: Take 02 misoprostol pills in every 3 hours interval until pregnancy passes.**